







MUSEUM
CARLSONIANUM.

MUSEUM CARLSONIANUM,

In quo

Novas et Selectas Aves,

Coloribus ad vivum brevique descriptione illustratas,

Suasum et sumptibus Generosissimi Possessoris;

Exhibet

ANDREAS SPARRMAN,

M. D. et Profess. Reg. Acad. Scient. Stockholmenf. Musei Praefect. Ejusd. Acad. ut et
Societ. Physiograph. Lund. Scient. ac Litt. Gothoburg. Hess. Homburg. Membr.



FASCICULUS II.



HOLMIAE,

Ex Typographia Regia, MDCCLXXXVII.

87
50.2
CHRB

Elenchus Avium Fasciculi II.

- N:o XXVI. *FALCO dubius.* XXXIX. *ANAS spectabilis*, Mas.
XXVII. *PSITTACUS cyaneus.* XL. — — — Femina.
XXVIII. — *Novae Zelandiae.* XLI. *LOXIA Cardinalis*,
varietas.
XXIX. — *Vernalis.* XLII. *EMBERIZA cyanella*,
Mas.
XXX. — *bimaculatus.*
XXXI. *ORIOIUS melaleucus.* XLIII. — — — Femina.
XLIV. — — — *ruticapilla.*
XXXII. *CUCULUS taitensis.* XLV. *TANAGRA capensis.*
XXXIII. *SITTA Chloris.* XLVI. *MUSCICAPA bicolor.*
XXXIV. *CERTHIA undulata.* XLVII. — — — *fuliginosa.*
XXXV. — — — *lepida.* XLVIII. *PARUS peregrinus*,
Mas.
XXXVI. — — — *armillata.*
XXXVII. *ANAS magellanica.* XLIX. — — — Femina.
XXXVIII. — *fraenata.* L. — — — *indicus.*
-



Falco dubius.

N:o XXVI.

FALCO dubius: corporis totius plumis ad
exortum albis, macula subalari magna, albida.

Rostrum nigricans.

Cera fecundum aetatem a flavo ad obscurum variat,
ut in aliquot Falconis speciebus non semel
compertum est.

Iris Lutea.

Pedes crocei.

Caput, *Colli Dorsum*, *Tectrices Alarum* cinereo-fuscae,
Plumarum singularum extrema margine f. acie fer-
ruginea.

Interscapulum, *Ipsumque Dorsum*, saturatius fusca.

Gula, *Jugulum*, *Pectus*, ferruginea, plumarum rachi-
dibus nigris.

Cauda superne cinerea, apice nigricans, versus basin
fasciis tribus fuscis notata.

—— subtus cinereo-albicans, apice fusca, pennis
fascia una alterave notatis.

OBS. Hanc speciem, quae in Svecia passim, sed non frequen-
ter occurrit, et pro Gyrfalconisvari etate haud merito a non-
nullis suspecta est, *F. dubium* appellavimus.

DEI



Psittacus cyaneus.

N:o XXVII.

PSITTACUS cyaneus : corpore toto
saturate et splendide Coeruleo.

Rostrum pallidum , apicē fuscum.

Pedes nigri.

Corpus totum ex cyaneo nitens.

OBS. Habitat in Infula Taiti f. Otaheiti. In hac quoque Cythera Maris Pacifici , ejusdem Psittaci varietatem pectore albo, illustrissimo Buffonio Taitensis dictam, (confr. *Planches enl.* 455. *fig.* 2.) soepe vidimus eaque Museum Carlsonianum ditavimus. Colorem autem cyaneum in utraque varietate eundem et hinc l. c. nimis dilutum esse monendum est.



Psittacus Novae Zelandiae.

N:o XXVIII.

PSITTACUS Novae Zelandiae: viridis, sincipite, macula sub-oculari et Hypochondriis coccineis.

Rostrum nigrum.

Frons, Sinciput, Macula utrinque sub-ocularis coccinea.

Occiput, Colli dorsum, Dorsum ipsum, Tectrices Alarum, Uropygium Caudaeque pagina superior prafina.

Jugulum, Pectus, Abdomen, Femora laete viridia.

Hypochondria maculis coccineis notata.

Alae et Cauda subtus fuscae.

Remigum Quinque Primariae fuscae, Disco anteriore superius dilatato et caerulescente.

Pedes nigri.

OBS. Duarum specierum hujus Generis, quae in Nova Zelandia occurrunt, haec circiter dupplo minor erat.



Psittacus vernalis.

N:o XXIX.

PSITTACUS vernalis : laete viridis,
alis prasinis, uropygio caudaeque pagina supe-
riore sanguineis, rostro pedibusque pallidis.

Rostrum rubicundum.

Pedes pallidi.

Tectrices Alarum saturatissime virides.

Remiges praeinae.

Uropygium sanguineum.

Cauda superius sanguinea.

—— *inferius* caerulea.

Reliqua Corporis pallide viridia.



Psittacus bimaculatus.

N:o XXX.

PSITTACUS bimaculatus : maculis
colli duabus oblongis , nigris , utriusque alae
macula magna et sulphurea.

Rostrum sub - fanguineum.

Frons , *Gula* , *Jugulum* , *Latera Capitis* Collique sub-
aurantia.

Gula et *Jugulum* a rostro ad initium pectoris *Macula*
Magna Ovali Nigra ex congerie macularum mino-
rum utrinque notantur.

Nucha , *Dorsum Colli* , *Interscapulium* , *Dorsum Ipsum* , *Uro-*
pygium *Caudaeque* pagina superior prafina.

Pectus , *Abdomen* , *Criſsum* laete viridia.

Tectrices Antiores *Minores Remigum Primar.* prafinae.

———— *Inferiores Majores Remigum Secundariarum* et
Remiges Posticae *Minores* glaucae , margine sulphureae.

Remigum Extima tota nigra.

———— *Reliquae Rachide* nigrae , *Disco* *exteriore* viri-
des , *Ultimo* *margine* leviter flavicantes , *Disco in-*
teriore fuscae ac *Ipsa Acie* sulphureo flavicantes.

Pedes fusco - cinerei.

Cauda subtus pallida.

Handwritten text, likely bleed-through from the reverse side of the page. The text is mostly illegible due to fading and bleed-through, but appears to be organized into several lines or paragraphs. Some faint words like "The" and "and" are visible.



Oriolus melaleucus.

N:o XXXI.

ORIO LUS melaleucus : *niger* , *Alarum*
macula alba.

Rostrum supra nigrum, infra fuscum.

Caput , *Collum* *Dorsumque* ex nigro nitentia.

Tectrices supra - et sub - alares minores albae.

—— *Reliquae* fuliginosae.

Remiges circiter viginti fuliginosae.

Pedes nigricantes.

Cauda rotundata Duodecim Rectricibus.

OBS. E Surinam pro ornando Museo Carlsoniano , Melaleuci
nostri specimen missum est.



Cuculus taitensis

N:o XXXII.

CUCULUS taitensis: *supra fuscus alboque punctatus, subtus laete ferrugineus ac lineolis brevibus longitudinaliter nigro striatus, cauda fusca maculis albidis sublunatis fasciata.*

Rostrum fusco flavicans.

Pedes nigri.

Capitulum totum nigricans, maculis oblongis testaceis ex rachidibus plumarum ejusdem coloris, notatum.

Colli latera, Dorsum Colli ac Interscapulum fusca maculisque oblongis albidis adspersa.

Alae, Dorsum, Uropygium maculis s. punctis majoribus albidis notata.

Gula, Jugulum, Pectus, Abdomen, Crissum, Femora, laete ferruginea, lineolis s. striis nigris longitudinalibus notata.

Rectrices fusco-nigrae, rachidibus brunneis, maculis untrunque ad discum sub-lunatis albidis fasciatae.



Sitta cloris.

N:o XXXIII.

SITTA Chloris: corpore supra viridi, subtus candido, cauda atra, ultimo apice flavicante.

Rostrum porrectum capite longius, subtus versus apicem nigricans.

Caput, *Dorsum Colli*, *Interscapulum Dorsumque* viridia.

Gula, *Jugulum*, *Pectus*, *Abdomen Crissumque* alba.

Remiges fuscae, *Margine* exteriori virescentes, transversaliter in Medietate flavicantes, hinc *Fascia* alarum transversalis tenuis flavescens.

Tectrices Alarum superiores f. minores virescenti-fuscae.

———— Inferiores anteriores, ac Superiores anteriores in angulo ad flexuram candidae.

Uropygium flavicans.

Cauda brevis, *Rectricibus* atris, extremo apice flavicantibus.

OBS. In Provincia Promontorii Bonae Spei, AKTER BRUNTIES HOOGTENS dicta, hanc speciem detegimus.



Certhia undulata.



N:o XXXIV.

CERTHIA undulata : *supra cinereo-fuliginosa* , *subtus alba nigroque transversim undulata*.

Rostrum arcuatum , *subtrigonum* , *fuscum*.

Capitium , *Latera Capitis* , *Dorsum Colli* , *Dorsum Ipsum* ,
Alae Caudaeque pagina superior cinereo-fuliginosa.

Gula , *Jugulum* , *Pectus* , *Abdomen* , *Femora ac Crissum*
alba , *lineis tenuibus transversis nigris undulatis fasciata*.

Remiges fuliginosae , *marginibus cinero-olivaceis*.

Pedes nigri.

Caudae pagina inferior cinerea.



Certhia lepidula.

N:o XXXV.

CERTHIA lepida: *subtus flava, pileo ex viridi nitente, collo et interscapulio hepaticis, uropygio splendide violaceo, alis, cauda pedibusque fuscis.*

Rostrum arcuatum, subtriquetrum, nigrum.

Capitulum viride nitens.

Capitis latera, Collum, Interscapulum intense hepatica.

Rectricum Alarum Una Alterave supremarum ad angulum alae interdum ex viridi nitent.

————— *Reliquae hepaticae.*

Remiges Fusco - Brunnei.

Dorsi Pars Posterior et Uropygium ex violaceo nitentia.

Pectus, Abdomen, Crissum, Femora citrina.

Pedes nigri.

Cauda nigra, Rectricibus flavis.



Certhia Armilata.



N:o XXXVI.

CERTHIA armillata : *viridis* , *alis complicatis supra nigris* , *subtus luteis* , *humeris* , *femorum armillis maculisque aliquot uropygii saphirinis*.

Rostrum arcuatum nigrum.

Capitulum , *Dorsum Colli* , *Dorsum Ipsum* smaragdino-viridia.

Gula , *Jugulum* , *Pectus* , *Tempora* , *Abdomen* , *Hypochondria* albido-virentia.

Criusum flavicans.

Alae Complicatae supra atrae , angulo exteriori f. *humeris* macula obliqua ex caeruleo - saphirino nitenti notatae , *subtus luteae* , margine exteriori et apice nigrae.

Remiges circiter viginti , marginibus exterioribus et apicibus nigris , marginibus autem interioribus flavis , ita ut in pennis complicatis luteum ex nigredine , nigredo autem ex luteo cooperiatur.

Pedes flavicantes ungulis nigris.

Cauda Rectricibus Duodecim constans.

OBS. Habitat in Surinam. Interdum variat plumula una alterave regionis iliaci , pectoris , extimi crissi ac temporum caeruleo-coelestibus. Taenia nigra a rostro per oculum ducta et hujus parti superiori contigua linea flava non raro variat.



Unas magellanica.



N:o XXXVII.

ANAS magellanica, *Femina*: *Speculo*
alarum viridi, *abdomine*, *crisso*, *uropygio*,
femoribus caudaque albis.

Rostrum cinereo-flavicans.

Capillitium cinereum.

Latera Capitis, *Gula* et *Colli pars superior* fusco-grisea,
Interscapulum, *Dorsum*, *Uropygium* fusco-brunnea.

Juguli, *Pectoris*, et *Abdominis Plumae* nigrae, fasciis
quatuor albis notatae.

Remiges Primariae nigrae.

———— *secundariae* albae.

Tectrices supra alares anteriores albae.

———— ————— *posteriores* ex viridi anatino tinctae
speculum formant.

Cauda Crissum et *Femora* alba.

Obs. Mās totus est albus, rostro nigro pedibusque flavis ac
Anserē domestico paulo minor; (cfr. celeberrimi Forsteri
Voyage round the world. Vol. II. pag. 495. *Femina* autem
Anseri magellanico Buffonii pl. enl. 1006. quodammodo
similis, ab hoc in plurimis tamen differt.

*Anas fraenata*



N:o XXXVIII.

ANAS fraenata: fusco - ferruginea, speculo alarum, abdomine capitisque annulo ad rostri basin albis.

Rostrum nigrum latiusculum.

Caput brunneum, ad *Rostri basin* *Habena* s. *Fascia* alba cinctum.

Collum ferrugineum.

Interscapulum, *Dorsum*, *Cauda* fusco - ferruginea.

Humeri, *Latera Pectoris* et *Hypochondria* punctis minutis albidis adspersa.

Pectus anteriùs superiùs ex fusco alboque undulatum.

Abdomen anteriùs albo - fericeum, posteriùs ex fusco fordide album.

Remiges Primariae Decem, nigrae.

—— *secundariae* albae, basi *Tectricibus* nigris coopertae; hinc speculum alarum parvum album.

Pedes nigri.

Cauda rotundata.

OBS. *Anatis fraenatae* specimen e Museo suo benigne communicavit Generosissimus ALSTRÖMER. Circa Alandiam occurrit et a venatoribus nonnullis pro *Anatis Marilae* femina habetur?



Anas spectabilis, Mas.



N:o XXXIX.

ANAS spectabilis, Mas : rostro basi gibbo
compresso : carina pennacea nigra capite ca-
nescente.

Rostri Gibbus flavus *Carina* pennacea nigra.

—— *ipsius* color ex rubicundo nigroque fuscus.

Capitulum ex plumbeo caerulescentique canum.

Oculi macula parva nigra cincti.

Regio infra-ocularis leniter virescens.

Gulae Nigredo versus nucham bifurcata extenditur.

Jugulum ac *Pectus* flavescenti-albida.

Colli Dorsum, *Humeri*, *Interscapulum*, alba.

Uropygium nigrum, *Lateribus* album.

Tectrices supra-alares anteriores inferiores candidae hu-
meros albos formant.

—— reliquae nigrae, earumque utriusque alae una
erecta et cochleata.

Remiges omnes nigricantes.

—— secundariae antrorsum falcatae.

Pedes rubicundi.

Cauda brevis, fuliginosa.

Obs. Edvardi figura, a Linneo citata, naturae minus congrua.



Anas spectabilis, Foemina.



N:o XL.

ANAS spectabilis, *Femina*: ferrugineo-
fusca nigroque maculata, regione oculorum alba,
apicibus alarum caudaque cinereis.

Rostrum nigrum.

Oculi macula parva alba cincti.

Caput et *Collum* ferruginea.

Pectus, *Interscapulium*, *Dorsum*, *Uropygium* ferruginea
nigroque maculata.

Abdomen, *Femora*, *Crissum* ferrugineo-fuliginosa.

Tectrices Omnes cinereo fuscae.

—— *Posteriores majores* apicibus albis, unde linea
transversalis alaris alba anterior formatur.

Remiges Primariae apice cinerae.

—— *Secundariae* nigricantes, apicibus obscure al-
bicantibus, hinc linea transversalis alaris albida
posterior.

Pedes nigri.

Cauda cinerea.

Obs. Qui artem venatoriam in mari Balthico ad oras Suderman-
niae exercent, hanc non nisi pro *Anatis spectabilis* Femina
habent.



Loxia Cardinalis



N:o XLI.

LOXIA Cardinalis, Varietas non cristata :
rubra, mento nigro, uropygio, cauda, alis
pedibusque fuscis.

Rostrum sub sanguineum.

Capite non cristato *Fronteque* minime nigra a varietate cristata praecipue differt.

Menti macula parva nigra differt a varietatis cristatae macula magna nigra quae mentum totamque gulam occupat.

Pedibus, *Alis*, *Caudaque* Varietati cristatae nec non *Loxiae mexicanae* Brissonii parum dissimilis, ab hac tamen magnitudine mentique nigredine praecipue differt.

Obs. Vivam ex Infulis Indiae occidentalis missam observavimus et ob habitum *Loxiae Cardinali* simillimum non nisi pro ejusdem pullo vel varietate habendam duximus.



Emberitza cyanella, Mas.



N:º XLII.

EMBERIZA cyanella, Mas: capite, collo, uropygio corporeque subtus caeruleo nitentibus; interscapulio humerisque ex caeruleo ferrugineoque variis, remigibus rectricibusque fuscis margine caerulefcentibus.

Rostrum pallide flavicans.


Caput, *Collum*, *Pectus*, *Abdomen*, *Femora*, *Crissum* et *Uropygium* cyanea.

Interscapulii, *Dorsi* *Tectricumque Plumae* cyanae, ferrugineis caeruleisve plumis margine apiceque rubiginosis mixtae.

Remiges, *Rectricesque* fuliginoso-fuscae marginibus caerulefcentibus.

Pedes flavicantes.

Obs. Habitat in America Septentrionali.





Emberitza cyanella, Foemina.



N:o XLIII.

EMBERIZA cyanella, *Femina*: corpore supra ex fusco, ferrugineo ac paululum caerulefcenti vario; subtus ferrugineo - albicante, maculis cyaneis, notato; remigibus rectricibusque fuscis, margine caerulefcentibus.

Rostrum pallide flavicans.

Capitulum, *Dorsum Colli*, *Interfcapulium*, *Dorsum Ipsum* ferruginea, plumis aliquot caeruleis hinc illinc sparsis variegata.

Gula albicans cyaneo maculata.

Jugulum, *Humeri* et *Pectus* ferruginea, maculis cyaneis dense notata.

Abdomen albicans caeruleo maculatum.

Crissum laete ferrugineum.

Rectrices Supra-Alares ac *Caudae* in medio caeruleae, ambitu rubiginosae.

—— *Sub-Alares* ac *Hypochondria* laeniter-virescentia.

Remiges Rectricesque fuscae margine cyaneae.

Pedes flavicantes.



Emberitza ruticapilla



N:o XLIV.

EMBERIZA ruticapilla : corpore supra
fusco, subtus cinereo; gula ferruginea, capistro
albo, capilitio insuper rufescente, cauda nigra.

Rostrum fusco-brunneum.

Capistrum ex macula magna albida nigro punctata,
frontem et genas occupante.

Vertex et *Occiput* rubicunda, anterieus et lateribus
nigro marginata.

Latera Colli, *Interseapulum*, *Alae* ac *Dorsum* fordide
fusca.

Gula ferruginea.

Pectus cinereo-ferrugineum.

Abdomen *Criussumque* cinerea.

Pedes fusco-brunnei.

Cauda fuliginoso-nigricans.



Tanager capensis.



N:o XLV.

TANAGRA capensis : *supra ferrugineo-fusca, subtus ferrugineo alboque varia, cauda in medio nigricante lateribus ferrugineo-rufescente, rostro flavicante, pedibus nigris.*

Rostrum pallide flavicans.

Frons ferruginea.

Vertex, Occiput, Latera et Dorsum Colli, Alae, Inter-
scapulum ac Dorsum ferrugineo-nigricantia.

Gula, Jugulum, Pectus, Abdomen, Femora ac Crissum
ferrugineo alboque varia ex plumis albis ferru-
ginoso marginatis.

Caudae medium nigricans, latera ferrugineo-rufescentia.
Pedes nigri.



Muscivora bicolor



N:o XLVI.

MUSCICAPA bicolor, corpore supra
cinereo, subtus ochraceo, cauda ex rectricibus
ochraceis cinereisque.

Rostrum nigricans.

Capitulum, *Dorsum Colli*, *Humeri*, *Interscapulum* ac *Dorsum* cinerea.

Rectrices Alares Inferiores cinerae, margine canescentes.

Remiges cinerae.

Pedes nigricantes.

Cauda longitudine circiter corporis et *Rectricibus* ochraceis cinereisque constans.

Obs. Habitat in Africa Meridionali.





Muscivora fuliginosa



N:o XLVII.

MUSCICAPA fuliginosa: corpore ex
ferrugineo et nigricante fuliginoso, Alis cauda-
que nigricantibus, rostro flavicante.

Rostrum pallide flavicans.


Caput, *Collum*, *Corpus supra* ac *subtus* nigricantia.

Alae nigricant.

Pedes nigri.

Cauda nigra, corporis mensura et dimidii.

OBS. In Deferto Africano inter rivulum Heuj et Fontem Quam-
medacka deteximus.





Larus peregrinus, Mas.



N:o XLVIII.

PARUS perigrinus, Mas: capite, collo dorsoque cinereis, pectore, macula alari uropygio ac rectricibus lateralibus postice coccineis.

Rostrum nigrum.

Lorum *Maculaque* sub-ocularis nigricant.

Capitulum, *Gula*, *Colli dorsum* ac *latera*, *Interscapulum*.
Dorsumque plumbeo-cinerea.

Rectrices Sub-Alares fulphureae.

—— *Supra-Alares* atrae, quarum *Aliquot Intermediae* apice coccineo-fulvae.

Remiges Primariae Duodecim, harum tres vel quatuor primae fusco-nigrae, margine interiore in medio paululum flavo tinctae.

—— *Primariae reliquae* ac

—— *Secundariae* fere omnes atrae in medio coccineae.

Juguli pars, *Pectus*, *Uropygium* coccinea.

Abdomen Crissumque leniter flavicantia.

Pedes nigri.

Cauda Rectricibus Decem nigris, tribus utrinque postice coccineis.

Obs. Character *Motacillae cinnamomeae* (confer. Syst. Nat. Edit. XII. pag. 335. N:o 32.) cum *P. peregr.* nostr. satis convenit, in variis tamen differt.



Larus peregrinus, foemina .



N:o XLIX.

PARUS perigrinus, Femina: corpore supra cinereo, subtus albo, macula alari aurantia, uropygio et rectricibus lateralibus postice coccineis.

Rostrum nigrum.

Capitulum, *Colli Dorsum*, *Interscapulum Dorsumque* cinerea.

Uropygium anterius flavo-virescens, posterius coccineo-fulvum.

Gula, *Jugulum*, *Pectus*, *Abdomen* fordide alba.

Cristum flavicans.

Tectrices Supra-Alares anteriores nigrae.

———— Inferiores posteriores flavae.

Remiges Primariae anteriores tres vel quatuor fuscae.

———— Reliquae fere omnes nigrae, basi fulvo-coccineae, una cum tectricibus posterioribus maculam alarem constituentes.

Pedes nigri.

Cauda Rectricibus ut in Mare, sed quinta utrinque etiam nigra apice oblique coccinea.

Obs. Confer. Syst. Nat. pag. 342.





Larus indicus.



N:o L.

PARUS indicus : gula et jugulo sordide
albis , pectore abdomine crissoque ferrugineis ,
corpore supra cinereo , rostro , pedibus , alis cau-
daque nigro-fuscis.

Rostrum fuscum.

Sinciput cinereo - nigricans.

Oculorum regiones , *Occiput* , *Colli* *Dorsum* , *Interscapulum*
Dorsum ac *Uropygium* cinerea.

Gula ac *Jugulum* sordide albicantia.

Pectus , *Abdomen* , *Crissum* ferruginea.

Tectrices Supra-Alares nigrae , margine cinerae.

Remiges fuliginoso - nigricantes.

Cauda supra fusco - nigricans , subtus cinereo - fusca.







SMITHSONIAN INSTITUTION LIBRARIES



3 9088 01203 8972